

Unravelling Entanglement

André R.R. Carvalho, Marc Busse, Olivier Brodier, Carlos Viviescas, and Andreas Buchleitner
Max-Planck-Institut für Physik komplexer Systeme, Nöthnitzer Strasse 38, D-01187 Dresden
 (Dated: February 1, 2008)

We show that the time evolution of entanglement under incoherent environment coupling can be faithfully recovered by monitoring the system according to a suitable measurement scheme.

PACS numbers: 03.67.-a, 03.67.Mn, 03.65.Yz, 42.50.Lc

Quantum entanglement, arguably the most non-classical feature of quantum mechanics, and considered as one of the key resources for quantum information processing, remains a puzzle for our intuition. General characterizations of the static entanglement and/or the separability properties of a given quantum state refer to the intricate geometry of tensor spaces, or to the operation of linear operations thereon. No general observable is known so far which would complement such essentially mathematical notions with a specific experimental measurement setup. Here we come up with a radically different, dynamical characterization of entanglement, through the continuous observation of a quantum system which evolves under incoherent coupling to an environment. We show that there is an optimal measurement strategy which provides an optimal decomposition of the time evolved, mixed system state, in the following sense: The entanglement of the time evolved mixed state is given by the average over the pure state entanglement of the single realisations of the stochastic time evolution. The weight of the different pure states is herein determined by their relative abundance during the detection process. Since arbitrary measurement prescriptions in general do *not* generate mixed state decompositions which are optimal in this sense, our finding implies that there are specific system observables which filter out the minimal amount of nonclassical correlations inscribed into a dynamically evolved, mixed quantum state.

Consider a bipartite quantum system composed of subsystems A and B , interacting with its environment. Due to this coupling, an initial pure state $|\Psi_0\rangle$ of the composite system will evolve into a mixed state $\rho(t)$, in a way governed by the master equation $\dot{\rho} = \sum_k \mathcal{L}_k \rho$. The superoperators $\mathcal{L}_k$ describe the effects of the environment on the system, and usually take the form [1]

$$\mathcal{L}_k \rho = \frac{\Gamma_k}{2} \left(2 J_k \rho J_k^\dagger - J_k^\dagger J_k \rho - \rho J_k^\dagger J_k \right), \quad (1)$$

where the operators J_k depend on the specific physical situation under study.

To extract the time evolution of entanglement under this incoherent dynamics, one solution is to evaluate a given entanglement measure $M(\rho)$ for the solution $\rho(t)$, at all times t . One starts from one of the known pure state measures $M(\Psi)$ [2, 3], together with a pure state

decomposition of ρ ,

$$\rho = \sum_i p_i |\Psi_i\rangle\langle\Psi_i|, \quad (2)$$

where the p_i are the positive, normalized weights of each pure state $|\Psi_i\rangle$. The most naive generalization for a mixed state would then be to consider the average

$$\overline{M} = \sum_i p_i M(\Psi_i), \quad (3)$$

which, however, is not uniquely defined, since the decomposition (2) is not unique. Therefore, the proper definition of $M(\rho)$ is the infimum of all possible averages $\overline{M}$ [4]. Finding this infimum has two main drawbacks: (i) it turns into a hard numerical problem for higher dimensional or multipartite systems, and, (ii) even for bipartite qubits, where analytical solutions for some measures $M(\rho)$ are known, there is no obvious general interpretation of the optimal decomposition, which gives the infimum, in physical terms.

Our approach here will be to avoid the direct use of the mixed state solutions $\rho(t)$, by substituting them by physically motivated ensembles of pure states. To do this, instead of solving the evolution equation for the density operator, we will follow a stochastic time evolution [5, 6, 7] of the initially pure state. This combines randomly occurring quantum jumps, defined by the action of the operators J_k in Eq. (1), with periods of continuous evolution, generated by a non-hermitian effective Hamiltonian. The resulting quantum trajectories, see Fig. 1, are known to provide, after averaging over many independent realisations, the same result as the master equation. Moreover, a single trajectory can be understood as the experimentally accessible information through the continuous monitoring of the system [6, 7, 8, 9]. In other words, one possible decomposition of the form (2) is given by the set of the possible outcomes $|\Psi_i\rangle$ of different runs of one and the same experiment, after a monitoring time t .

Note, however, that the stochastic time evolution induced by the jump operators J_k does not necessarily yield the optimal decomposition which minimises $\overline{M}$. Yet, also the choice of the jump operators is not unique: Indeed, the definition of new jumps as linear combinations of the original ones, $\tilde{J}_k = \sum_i U_{ki} J_i$, with U a unitary matrix, or by the addition of a complex number μ ,




FIG. 1: Simplified schematic representation of stochastic time evolutions generated by continuous monitoring of the system dynamics through a jump operator J : During each time step of (short) duration δt , the system evolves with a certain probability δp into the state $|\Psi_{\delta t}^1\rangle$, and with probability $(1 - \delta p)$ into the state $|\Psi_{\delta t}^2\rangle$, depending on the result of the action of J on the initial state $|\Psi_0\rangle$. Upon iterative application of this method, a large number of different states can be accessed, giving rise to different stochastic pure state trajectories. At a given time $t = n\delta t$, this process generates a pure state decomposition of the time evolved density matrix $\rho(t)$.

$J'_{k,\pm} = (\mu \pm J_k) / \sqrt{2}$, leaves Eq. (1) unaltered. More generally, one can perform both changes simultaneously, to obtain the general jump operators $L_{k,\pm} = (\mu \pm \tilde{J}_k) / \sqrt{2}$.

Since each such choice of jump operators produces a different *unravelling* of the same master equation (1), i.e., a different decomposition of $\rho(t)$, we have to search for that type of monitoring which reproduces the desired optimal decomposition, upon averaging over the different realisations at time t . We claim that such a measurement prescription can always be found, for a given initial state, and given environment coupling. While a rigorous mathematical proof of this conjecture is still missing, we now present strong evidence in its favour, for various decoherence models.

We focus on the dynamics of bipartite two-level systems, where the results obtained by our method can readily be compared with known analytical solutions [3, 10]. We assume that each subsystem interacts independently with its own environment, and consider those situations where decoherence is induced by dissipation (zero temperature reservoir), by dephasing, or by noise (infinite temperature reservoir). For simplicity, we set all decay rates $\Gamma_k \equiv \Gamma$ in Eq. (1). We start with the general jump operators $L_{k,\pm}$ as defined above, and search for the ones which yield the optimal solution. The first step in such procedure is to follow all possible states $|\Psi_{\delta t}\rangle$ of the system, at a short time δt after the initial time $t_0 = 0$, and to calculate the average measure $\overline{M}(\delta t)$. Since the possible states $|\Psi_{\delta t}^k\rangle$ which can be reached after this short time interval are given by the action of the N jump operators, plus the state $|\Psi_{\delta t}^{N+1}\rangle$ which results from continuous evolution (see figure 1), the expression for the average measure is simply $\overline{M}(\delta t) = (1 - \sum_{k=1}^N \delta p_k) M(\Psi_{\delta t}^{N+1}) + \sum_{k=1}^N \delta p_k M(\Psi_{\delta t}^k)$,




FIG. 2: Entanglement dynamics of a bipartite quantum state initially prepared in the maximally entangled state $|\Psi_0\rangle = (|00\rangle + |11\rangle) / \sqrt{2}$, under incoherent coupling to a zero temperature environment. $\overline{M}$ is here derived from the entanglement of formation [2]. The symbols result from an average over different realisations of the stochastic time evolution under the action of the general jump operators $L_{k,\pm}$, while the solid line represents the exact result derived from the integration of the master equation. Open squares ($\mu = 1$) and filled circles ($\mu = 3$) are obtained by progressively improved unravellings, with an optimal choice of the unitary U . Open triangles indicate the result of an arbitrarily chosen unravelling, with no optimization, neither of U nor of μ .

and can be minimised over the different possible unravellings. This ensures that the short time behaviour of the average measure is optimal.

In fact, for many initial states, such as the Bell states $(|00\rangle + |11\rangle) / \sqrt{2}$ and $(|01\rangle + |10\rangle) / \sqrt{2}$ under dephasing, or a state $|\Psi_0\rangle = \psi_{00}|00\rangle + \psi_{01}|01\rangle + \psi_{10}|10\rangle$ coupled to a zero temperature environment, the minimisation for the initial step can be shown to be not only a necessary but also a sufficient condition, for $\overline{M}(t)$ to coincide with $M(\rho(t))$. This is a striking feature, since the optimisation needs to be performed only once, for the initial time step, but leads to a faithful, optimal time evolution for arbitrary $t > t_0$.

In other cases, however, the correct behaviour at short times is not sufficient, and further search for the optimal unravelling is needed, though now in a reduced parameter space (spanned by the parameters of $L_{k,\pm}$). Such search is illustrated in Fig. 2, for a Bell initial state $|\Psi_0\rangle = (|00\rangle + |11\rangle) / \sqrt{2}$, coupled to a zero temperature reservoir, and for $M(\Psi)$ given by the familiar entanglement of formation [2]. The condition for the correct initial decay fixes most of the parameters of the unitary matrix U , and enforces $\mu \geq 1/\sqrt{2}$. The open squares and filled circles correspond to unravellings satisfying these conditions for $\mu = 1$, and $\mu = 3$, respectively, while triangles are obtained by a non-optimal choice of U and μ .




FIG. 3: Time evolution of the stochastically generated entanglement of formation (which we use here for the explicit evaluation of $\bar{M}$), during the application of a CNOT gate in a two qubit system coupled to a dephasing environment. Adequate choice of the jump operators $L_{k,\pm}$ leads to perfect agreement (circles) with the exact result (solid line) during the gate time. A non-optimal choice of unravelling leads to large deviations from the exact solution.

Although both values of μ induce the correct behaviour for sufficiently short times, only the larger value of μ gives good agreement for long times. The inset shows the situation for the same initial state, though coupled to an infinite temperature environment. The unravelling is chosen to give the correct initial slope, but the agreement is also very good for the asymptotics. The only discrepancy persists at intermediate times, where the exact solution for $M(\rho(t))$ exhibits a discontinuous derivative.

We can allow for more general dynamics, with a non-trivial unitary evolution in the presence of incoherent environment coupling. This is relevant when it comes to describe, for example, gate operations performed in open quantum systems. To illustrate such a situation, we simulated the action of a CNOT gate, in a system coupled to a dephasing reservoir as shown in Fig. 3. Starting with a separable initial state $|\Psi_0\rangle = (|00\rangle + |10\rangle)/\sqrt{2}$, the gate would generate the maximally entangled state $|\Psi\rangle = (|00\rangle + |11\rangle)/\sqrt{2}$, after a time T_{gate} , chosen to be five times faster than the decoherence time scale $1/\Gamma$ (see Eq. (1)). However, due to dephasing, the entanglement at T_{gate} will not be maximal, since the target state preparation will not be perfect. As in our above examples one can find an unravelling which generates an optimal pure state decomposition, in excellent agreement with the exact entanglement evolution obtained from the solution $\rho(t)$. Once again, a non-optimal unravelling suggests a very different entanglement dynamics, with apparently almost perfect gate performance – in pronounced contrast to the correct result.

Finally, let us consider the example of a tripartite qubit




FIG. 4: Multipartite entanglement decay for a GHZ state under dephasing dynamics. Filled circles represent the tripartite concurrence $C_3(t)$ (substituted in the explicit evaluation of $\bar{M}$) [11] obtained after averaging over 1000 realisations of the unravelling defined by the jump operators $J_k = \sigma_+^{(k)} \sigma_-^{(k)}$ ($k = 1, 2, 3$) (which actually coincide with the coupling operators in the dephasing master equation), in very good agreement with the exact solution indicated by the solid line. A non-optimal unravelling (open circles) considerably overestimates the actual mixed state entanglement of the time evolved state.

system, initially prepared in a GHZ or in a W state, and in contact with dephasing or zero temperature environments, respectively. In both cases, the optimal unravelling produces excellent agreement with the exact result for the tripartite concurrence $C_3(t)$ [3, 11], as illustrated in Fig. 4 for a GHZ initial state subject to dephasing. In this exemplary case, the optimal detection is provided by the original jump operators $J_k = \sigma_+^{(k)} \sigma_-^{(k)}$, with $k \in \{1, 2, 3\}$. Again, a non-optimal unravelling leads to very different predictions.

To conclude, we have shown that mixed state entanglement can be unraveled by a suitable measurement prescription. The latter can be derived from the initial state of the composite quantum system under study, together with the specific type of environment coupling. There is thus no ambiguity (which was suggested in [12]) in the definition of mixed state entanglement with respect to different unravellings: the minimal nonclassical correlations needed to characterize a dynamically evolving quantum state are identified by an optimal (in general not unique) monitoring prescription, and any in this sense non-optimal unravelling provides a non-optimal estimate thereof.

Surprisingly, in all examples considered, we so far always succeeded to come up with a *time-independent* optimal unravelling, i.e., the monitoring strategy does not need to be readapted during the incoherent time evolution. This suggests a kind of uniform convergence of the

stochastic average towards the exact solution (which follows from the evaluation of the convex roof of a given entanglement measure on the time evolved density matrix), for the optimal unravelling, and implies a subtle relation between the temporal evolution of entanglement and the ensemble of pure states which can be reached

during the single realisations generated by the optimal jump operators.

We thank Florian Mintert, Daniel Estève, Armin Uhlmann, and Robin Hudson for entertaining discussions and useful suggestions.

-
- [1] G. Lindblad, Math. Phys. **48**, 119 (1976).
 - [2] W. K. Wootters, Phys. Rev. Lett. **80**, 2245 (1998).
 - [3] F. Mintert, A. R. R. Carvalho, M. Kús, and A. Buchleitner, Phys. Rep. **415**, 207 (2005).
 - [4] A. Uhlmann, Phys. Rev. A **62**, 032307 (2000).
 - [5] H. Carmichael, *An open systems approach to quantum optics*, Lecture Notes in Physics m 18 (Springer-Verlag, Berlin, 1993).
 - [6] K. Mølmer, Y. Castin, and J. Dalibard, J. Opt. Soc. Am. B **10**, 524 (1993).
 - [7] K. Mølmer and Y. Castin, Quant. Semiclass. Opt. **8**, 49 (1996).
 - [8] T. Sauter, W. Neuhauser, R. Blatt, and P. Toschek, Phys. Rev. Lett. **57**, 1696 (1986).
 - [9] H. M. Wiseman and G. J. Milburn, Phys. Rev. A **47**, 1652 (1993).
 - [10] W. K. Wootters, Phys. Rev. Lett. **80**, 2245 (1998).
 - [11] A. R. R. Carvalho, F. Mintert, and A. Buchleitner, Phys. Rev. Lett. **93**, 230501 (2004).
 - [12] H. Nha and H. J. Carmichael, Phys. Rev. Lett. **93**, 120408 (2004).